

Lithobates catesbeianus



Taxon	Family / Order / Class / Phylum
<i>Lithobates catesbeianus</i> (Shaw, 1802)	Ranidae / Anura / Amphibia / Vertebrata

COMMON NAMES (English only)

American Bullfrog

SYNONYMS

Rana catesbeiana Shaw, 1802

SHORT DESCRIPTION

It is the largest North-American ranid. In its natural range, adults can reach at most a 184 mm snout-vent length (Louisiana), while in Europe (southwestern France), adults may exceed a 195 mm snout-vent length, with an average weight of approximately 430 grams. It is likely to feed on a wide range of prey: amphibians, fishes, small mammals, ducklings and small bird species, molluscs, crustaceans and insects. In southwestern France, the adult diet consists of a wide range of prey, with amphibians and insects most prominent.



BIOLOGY/ECOLOGY

Dispersal mechanisms

Introductions by humans. Natural spread of juveniles and adults is terrestrial or via rivers.

Reproduction

In southwestern France, the breeding period begins in May and lasts until early September, and tadpole development takes one to two years.

Known predators

In southwestern France, larval and young frog stages seem to undergo significant predation. Known predators include carnivorous fishes, in particular *Micropterus salmoides* introduced from North America, and probably some Mustelidae and Ardeidae. The main source of predation, in high-density areas, appears to be cannibalism.

Resistant stages (seeds, spores etc.)

None.

HABITAT

Native (EUNIS code)

C: Inland surface water habitats

Habitat occupied in invaded range (EUNIS code)

C: Inland surface water habitats, D: Mire, bog and fen habitats

Habitat requirements

In its native range, it usually lives in lakes or large ponds. Where introduced, it occupies any type of habitat that is lentic or with slowly moving water, especially if aquatic and bank vegetation are abundant.

DISTRIBUTION

Native Range

The American Bullfrog is autochthonous in the eastern part of North America, from Vera Cruz north to the Great Lakes and New Brunswick.

Known Introduced Range

It is currently established in the western part of North America, between California and southern British Columbia, as well as in several countries of South America, some islands of the Greater Antilles, Asia, Hawaii and Europe. In Europe, it has been introduced in Germany, Italy, United Kingdom, Spain, the Netherlands, Greece, Belgium and France.




Trend

Although it is presently illegal to import this species to the EU, it can still be ordered on the Internet and shipped worldwide. Farming could increase its distribution area by escapes.

MAP (European distribution)



Legend

	Known in country		Known in CGRS square		Known in sea
---	------------------	---	----------------------	--	--------------

INTRODUCTION PATHWAY

It has been introduced in many parts of the world, often for farming purposes but also as a predator of unwanted species like insect pests in Hawaii, and sometimes as a pet.

IMPACT

Ecosystem Impact

Where introduced, it has the ability to occupy a whole range of habitats and to feed on many species. A negative impact on native ranids has been notably stressed. American Bullfrogs carry an emerging pathogenic fungus, *Batrachochytrium dendrobatidis*, which has been implicated in global amphibian decline.

Human Health and Social Impact

Unknown.

Economic Impact

Not documented but the cost of eradication has been US\$29 000 in the UK.

MANAGEMENT

Prevention

Unknown.

Mechanical

Except for a limited area in the United Kingdom and France, no eradication strategy has been formulated in Europe. Only few control operations have been attempted (i.e. in Germany). In southwestern France, a long-term regional program was started in May 2003, conducted by Cistude Nature (environment protection

organization). It aims at the development of strategies and management tools (in particular egg collection, tadpoles trapping, and juvenile and adult shooting) before early 2007, to try to limit American Bullfrog populations, or even to eradicate them.

Chemical

Unknown.

Biological

Unknown.

REFERENCES

- Détaint M, Coïc C (2006) La Grenouille taureau *Rana catesbeiana* dans le sud-ouest de la France. Premiers résultats du programme de lutte. Bulletin de la Société Herpétologique de France 117:41-56
- Garner TWJ, Perkins MW, Govindarajulu P, Seglie D, Walker S, Cunningham AA, Fisher MC (2005) The emerging amphibian pathogen *Batrachochytrium dendrobatidis* globally infects introduced populations of the North American bullfrog, *Rana catesbeiana*. Biology Letters 2:455-459
- Pearl CA, Adams MJ, Bury RB, McCreary B (2004) Asymmetrical effects of introduced Bullfrogs (*Rana catesbeiana*) on native ranid frogs in Oregon. Copeia 2004:11-20

OTHER REFERENCES

- Arano B, Llorente G, García París M, Herrero P (1995) Species translocation menaces Iberian waterfrogs. Conservation Biology 9:196-198
- Boettger CR (1941) Der Versuch einer Ochsenfroschzucht in der Lüneburger Heide. Sitzungsberichte der Gesellschaft Naturforschender Freunde zu Berlin, pp 216-221
- Bruneau M, Magnin E (1980) Croissance, nutrition et reproduction des ouaouarons *Rana catesbeiana* Shaw (Amphibia Anura) des Laurentides au nord de Montréal. Canadian Journal of Zoology 58:175-183
- Cross CL (2002) *Rana catesbeiana* (American Bullfrog) diet. Herpetological Review 33:129-130
- Cunningham AA, Langton TES (1997) Disease risks associated with translocations of Amphibians into, out of and within Europe - a U.K. perspective. The Journal of the British Veterinary Zoological Society 3: 37-41
- Détaint M, Coïc C (2001) Invasion de la Grenouille taureau (*Rana catesbeiana*) en France: Synthèse bibliographique - Suivi 2000-2001 - Perspectives. Association Cistude Nature. Le Haillan (France): www.grenouilletaureau.net
- Détaint M, Coïc C (2003) La Grenouille taureau: *Rana catesbeiana* Shaw, 1802. Pp 154-156. In : Pascal M, Lorvelec O, Vigne JD, Keith P, Clergeau P (eds) Évolution holocène de la faune de Vertébrés de France: invasions et disparitions. Ministère de l'Écologie et du Développement Durable, Paris
- Frazer JFD (1964) Introduced species of amphibians and reptiles in mainland Britain. British Journal of Herpetology 3:145-150
- Ficetola GF, Coïc C, Detaint M, Berroneau M, Lorvelec O, Miaud C. Pattern of distribution of the American bullfrog *Rana catesbeiana* in Europe. Biological Invasions. In press
- Frost DR, Grant T, Faivovich J, Bain RH, Haas A, Haddad CFB, De Sá RO, Channing A, Wilkinson M, Donnellan SC, Raxworthy CJ, Campbell JA, Blotto BL, Moler P, Drewes RC, Nussbaum RA, Lynch JD, Green DM, Ward C, Wheeler WC (2006) The amphibian tree of life. Bulletin of the American Museum of Natural History 297:1-370
- García París M (1991) Primeros datos sobre *Rana catesbeiana* Shaw, 1802 (Anura : Ranidae) en España. Revista Española de Herpetología 5:89-92
- Govindarajulu P (2004) Introduced bullfrogs (*Rana catesbeiana*) in British Columbia: impacts on native Pacific treefrogs (*Hyla regilla*) and red-legged frogs (*Rana aurora*). Ph.D. thesis, University of Victoria, Canada
- Hammerson GA (1982) Bullfrog eliminating leopard frogs in Colorado? Herpetological Review 13:115-116
- Hays MP, Warner J (1985) *Rana catesbeiana* Food. Herpetological Review 16:109
- Hecnar SJ, M'Closkey RT (1997) Changes in the composition of a Ranid frog community following bullfrog extinction. American Midland Naturalist 137:145-150
- Kiesecker JM, Blaustein AR (1997a) Effects of introduced bullfrogs and smallmouth bass on microhabitat use, growth and survival of native red-legged frogs. Conservation Biology 12:776-787
- Kiesecker JM, Blaustein AR (1997b) Population differences in responses of red-legged frogs (*Rana aurora*) to introduced bullfrogs. Ecology 78:1752-1760
- Kupferberg SJ (1997) Bullfrog (*Rana catesbeiana*) invasion of a California river: the role of larval competition. Ecology 78:1736-1751

- Lanza B (1962) On the introduction of *Rana ridibunda* Pallas and *Rana catesbeiana* Shaw in Italy. *Copeia* 1962:642-643
- Laufer H (2004) Zum beutespektrum einer population von Ochsenfröschen (Amphibia: Anura: Ranidae) nördlich von Karlsruhe (Baden-Württemberg Deutschland). *Faunistische Abhandlungen, Dresden*. 25:139–150
- Laufer H, Waitzman M (2002) Der Ochsenfrosch (*Rana catesbeiana*) am nördlichen Oberrhein (Baden-Württemberg). *Herpetofauna* 24:5-14
- Mazzoni R, Cunningham AA, Daszak P, Apolo A, Perdomo E, Speranza G (2003) Emerging pathogen of wild amphibians in frogs (*Rana catesbeiana*) farmed for international trade. *Emerging Infectious Diseases* 9:995-998
- Moyle PB (1973) Effects of introduced bullfrogs, *Rana catesbeiana*, on the native frogs of the San Joaquin Valley, California. *Copeia* 1973:18-22
- Neveu A (1997) L'introduction d'espèces allochtones de grenouilles vertes en France, deux problèmes différents : celui de *R. catesbeiana* et celui des taxons non présents du complexe *esculenta*. *Bulletin Français de la Pêche et de la Pisciculture* 344/345:165-171
- Shaw G (1802) *General Zoology or Systematic Natural History*. Society for the study of Amphibians and Reptiles, London 3:106-108
- Stumpel AHP (1992) Successful reproduction of introduced bullfrogs *Rana catesbeiana* in north-western Europe: a potential threat to indigenous amphibians. *Biological Conservation* 60:61-62
- Thiesmeier B, Jäger O, Fritz U (1994) Erfolgreiche Reproduktion des Ochsenfrosches (*Rana catesbeiana*) im nördlichen Landkreis Böblingen (Baden-Württemberg). *Zeitschrift für Feldherpetologie* 1:169-176
- Touratier L (1992) Première apparition en France (région Aquitaine) d'une grenouille géante américaine: *Rana catesbeiana* en voie d'acclimatation. Intérêt zoologique et impact éventuel sur l'environnement. *Bulletin de la Société Vétérinaire Pratique de France* 76:219-228
- Wavrin de H (2000) Note sur la présence de la Grenouille Taureau (*Rana catesbeiana*) en Brabant Wallon. *Projet de nouvel atlas de Wallonie et Bruxelles*, Maison Liégeoise de l'Environnement, Liège, Belgique

Author: Olivier Lorvelec and Mathieu Détaint

Date Last Modified: November 20th, 2006